



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 277/36, 239/38, 285/12, A61K 31/425, 31/41, 31/505		A1	(11) International Publication Number: WO 94/20480
			(43) International Publication Date: 15 September 1994 (15.09.94)
(21) International Application Number: PCT/CA94/00135 (22) International Filing Date: 10 March 1994 (10.03.94) (30) Priority Data: 030,924 12 March 1993 (12.03.93) US (60) Parent Application or Grant (63) Related by Continuation US 030,924 (CIP) Filed on 12 March 1993 (12.03.93) (71) Applicant (for all designated States except US): MERCK FROSST CANADA INC. [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Quebec H9H 3L1 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): BLACK, Cameron, W. [CA/CA]; 22 Saddlewood Pointe-Claire, Quebec, Quebec H9R 5S6 (CA). GUAY, Daniel [CA/CA]; 67 Roger Maillet, Notre Dame de l'Île Perrot, Quebec J7V 7P2 (CA). LI, Chun-Sing [GB/CA]; 2 Libersan Crescent, Dollard Des Ormeaux, Quebec, H9A 2B5 (CA). PRASIT, Petpiboon [CA/CA]; 177 Argyle, Kirkland, Quebec H9H 5A6 (CA).		ROY, Patrick [CA/CA]; 404 Hurteau, #312, Dollard Des Ormeaux, Quebec H9G 1X4 (CA). (74) Agent: MURPHY, Kevin, P.; Swabey Ogilvy Renault, Suite 800, 1001 de Maisonneuve Boulevard West, Montreal, Quebec H3A 3C8 (CA). (81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TT, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: ALKANESULFONAMIDO-1-INDANONE DERIVATIVES AS INHIBITORS OF CYCLOOXYGENASE			
(57) Abstract			
Disclosed are compounds of formula (I) useful in the treatment of cyclooxygenase mediated diseases such as pain, fever and inflammation of a variety of conditions including rheumatic fever, symptoms associated with influenza or other viral infections, common cold, low back and neck pain, dysmenorrhea, headache, toothache, sprains and strains, myositis, neuralgia, synovitis, arthritis, including rheumatoid arthritis degenerative joint diseases (osteoarthritis), gout and ankylosing spondylitis, bursitis, burns, injuries.		<div style="text-align: center;"> $R^1-S(O)_2-N-H$ (I) </div>	
Atty. Docket No. 01379/1/US Serial No. 10/769,485 Timothy Maziarz Reference B7			

Best Available Copy

Best Available Copy

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

TITLE OF THE INVENTION

ALKANESULFONAMIDO-1-INDANONE DERIVATIVES AS
INHIBITORS OF CYCLOOXYGENASE

5 BACKGROUND OF THE INVENTION

This invention relates to compounds and pharmaceutical compositions for the treatment of cyclooxygenase mediated diseases and methods of treating thereof.


10 Non-steroidal, antiinflammatory drugs exert most of their antiinflammatory, analgesic and antipyretic activity and inhibit hormone-induced uterine contractions and certain types of cancer growth through inhibition of prostaglandin G/H synthase, also known as cyclooxygenase. Up until recently, only one form of cyclooxygenase had been characterized, this corresponding to cyclooxygenase-1 or the
15 constitutive enzyme, as originally identified in bovine seminal vesicles. Recently the gene for an inducible form of cyclooxygenase (cyclooxygenase-2) has been cloned, sequenced and characterized from chicken, murine and human sources. This enzyme is distinct from the cyclooxygenase-1 which has now also been cloned, sequenced and
20 characterized from sheep, murine and human sources. The second form of cyclooxygenase, cyclooxygenase-2, is rapidly and readily inducible by a number of agents including mitogens, endotoxin, hormones, cytokines and growth factors. As prostaglandins have physiological and pathological roles, we have concluded that the constitutive enzyme,
25 cyclooxygenase-1, is responsible, in large part, for endogenous basal release of prostaglandins and hence is important in their physiological functions such as the maintenance of gastrointestinal integrity and renal blood flow. In contrast, we have concluded that the inducible form, cyclooxygenase-2, is mainly responsible for the pathological effects of
30 prostaglandins where rapid induction of the enzyme would occur in response to such agents as inflammatory agents, hormones, growth factors, and cytokines. Thus, a selective inhibitor of cyclooxygenase-2 will have similar antiinflammatory, antipyretic and analgesic properties of a conventional non-steroidal antiinflammatory drug (NSAID), and in

- 2 -

addition would inhibit hormone-induced uterine contractions and have potential anti-cancer effects, but will have a diminished ability to induce some of the mechanism-based side effects. In particular, such a compound should have a reduced potential for gastrointestinal toxicity, a reduced potential for renal side effects, a reduced effect on bleeding times and a lessened ability to induce asthma attacks in aspirin-sensitive asthmatic subjects.

SUMMARY OF THE INVENTION

The invention encompasses compounds of Formula I



I

which are useful in the treatment of cyclooxygenase mediated diseases, in particular cyclooxygenase-2 mediated diseases.

The invention also encompasses pharmaceutical compositions for inhibiting cyclooxygenase and for treating cyclooxygenase mediated diseases as disclosed herein comprising a pharmaceutically acceptable carrier and a non-toxic therapeutically effective amount of compound of Formula I as described herein.


The invention also encompasses methods of inhibiting cyclooxygenase and treating cyclooxygenase mediated diseases comprising:

administration to a patient in need of such treatment of a non-toxic therapeutically effective amount of a compound of Formula I as disclosed herein.

- 3 -

DETAILED DESCRIPTION OF THE INVENTION

The invention encompasses compounds of Formula I



I

and pharmaceutically acceptable salts thereof wherein:

R¹ is selected from the group consisting of

- 15
- (a) C₁-7alkyl, and
 - (b) mono-, di-, tri-, tetra- and per-substituted C₁-7alkyl, wherein the substituent is fluoro;

X is O, S, or -CH₂-;

Ar is a mono or disubstituted aromatic ring, said ring selected from

- 20
- (a) a ring of 5 atoms containing one O, S or N atom and optionally, 1, 2 or 3 additional N atoms; and
 - (b) a ring of 6 atoms containing 2, 3 or 4 nitrogen atoms;

wherein the substituents are independently selected from

- 25
- (a) hydrogen,
 - (b) C₁-7alkyl,
 - (c) halogen, including F, Cl, Br, and I,
 - (d) OCH₃,
 - (e) SCH₃,
 - (f) CF₃,
 - (g) COCH₃ and
 - (h) S(O)₂CH₃.
- 30

For purposes of this specification alkyl is defined to include linear, branched or cyclic alkyl including methyl, ethyl, 2-propyl, t-

- 4 -

butyl, 1,1-dimethylethyl, cyclopropyl, cyclobutyl, cyclohexyl, and cyclohexyl. Similarly fluorosubstituted alkyl as defined above is defined to include fluoromethyl, trifluoromethyl, 2-fluoroethyl, perfluoroheptyl, perfluorocyclopropyl, and perfluoroheptyl.

5

One genus of this embodiment concerns compounds of Formula I wherein:

Ar is a mono- or disubstituted aromatic group wherein the aromatic group is selected from the group consisting of

10

- (1) furanyl,
- (2) diazinyl, triazinyl, tetrazinyl,
- (3) imidazolyl,
- (4) isooxazolyl,
- (5) isothiazolyl,
- (6) oxadiazolyl,
- (7) oxazolyl,
- (8) pyrazolyl,
- (9) pyrrolyl,
- (10) thiadiazolyl,
- (11) thiazolyl,
- (12) thienyl,
- (13) triazolyl, and
- (14) tetrazolyl,

15

20

wherein the substituents are independently selected from

25

- (a) hydrogen,
- (b) C₁₋₇alkyl,
- (c) halogen, including F, Cl, Br, and I,
- (d) OCH₃,
- (e) SCH₃, and
- (f) CF₃,
- (g) COCH₃, and
- (h) S(O)₂CH₃

30

- 5 -

In one class of this genus the invention concerns compounds of Formula I wherein,

R¹ is selected from the group consisting of

- 5 (a) methyl, ethyl, propyl, and butyl
 (b) mono, di or tri substituted C₁₋₄alkyl, wherein the substituent is fluoro;

X is O, S or CH₂

Ar is a mono or disubstituted aromatic group wherein the aromatic group is selected from the group consisting of

- 10 (1) 2-furanyl,
 (2) 3-furanyl,
 (3) 2-thienyl,
 (4) 3-thienyl,
 (5) 3-isooxazolyl,
15 (6) 4-isooxazolyl,
 (7) 5-isooxazolyl,
 (8) 3-isothiazolyl,
 (9) 4-isothiazolyl,
 (10) 5-isothiazolyl,
20 (11) 2-oxazolyl,
 (12) 4-oxazolyl,
 (13) 5-oxazolyl,
 (14) 2-thiazolyl,
 (15) 4-thiazolyl,
25 (16) 5-thiazolyl,
 (17) 1,2,3-thiadiazol-4-yl,
 (18) 1,2,3-thiadiazol-5-yl,
 (19) 1,2,4-thiadiazol-3-yl,
 (20) 1,2,4-thiadiazol-5-yl,
30 (21) 1,3,4-thiadiazol-2-yl,
 (22) 1,2,5-thiadiazol-3-yl,
 (23) 1,2,3-oxadiazol-4-yl,
 (24) 1,2,3-oxadiazol-5-yl,
 (25) 1,2,4-oxadiazol-3-yl,

- 6 -

- 5 (26) 1,2,4-oxadiazol-5-yl,
(27) 1,3,4-oxadiazol-2-yl,
(28) 1,2,5-oxadiazol-3-yl,
(29) pyrazol-4-yl,
(30) pyrazol-5-yl,
(31) 1,2,3-triazol-4-yl,
(32) 1,2,3-triazol-5-yl,
(33) 1,2,4-triazol-3-yl,
(34) 1,2,4-triazol-5-yl,
10 (35) 1,2-diazinyl,
(36) 1,3-diazinyl,
(37) 1,4-diazinyl,
(38) imidazol-2-yl,
(39) imidazol-4-yl,
15 (40) imidazol-5-yl,
(41) tetrazol-5-yl,

wherein the substituents are independently selected from

- (a) hydrogen,
(b) C₁-3alkyl,
20 (c) halogen, including F, Cl, Br, and I,
(d) OCH₃,
(e) SCH₃,
(f) CF₃,
(g) COCH₃, and
25 (h) S(O)₂CH₃

In a second embodiment, the invention encompasses pharmaceutical compositions for inhibiting cyclooxygenase and for treating cyclooxygenase mediated diseases as disclosed herein comprising a pharmaceutically acceptable carrier and a non-toxic therapeutically effective amount of compound of Formula I as described above.

Within this embodiment the invention encompasses pharmaceutical compositions for inhibiting cyclooxygenase-2 and for

- 7 -



treating cyclooxygenase-2 mediated diseases as disclosed herein comprising a pharmaceutically acceptable carrier and a non-toxic therapeutically effective amount of compound of Formula I as described above.

5 In a third embodiment, the invention encompasses a method of inhibiting cyclooxygenase and treating cyclooxygenase mediated diseases as disclosed herein comprising:
administration to a patient in need of such treatment of a non-toxic therapeutically effective amount of a compound of Formula I as
10 disclosed herein.


Within this embodiment the invention encompasses a method of inhibiting cyclooxygenase-2 and treating cyclooxygenase-2 mediated diseases as disclosed herein comprising:
administration to a patient in need of such treatment of a non-toxic
15 therapeutically effective amount of a compound of Formula I as disclosed herein.

As disclosed elsewhere in this specification in further detail, these diseases include pain, fever and inflammation of a variety of conditions including rheumatic fever, symptoms associated with
20 influenza or other viral infections, common cold, low back and neck pain, dysmenorrhea, headache, toothache, sprains and strains, myositis, neuralgia, synovitis, arthritis, including rheumatoid arthritis degenerative joint diseases (osteoarthritis), gout and ankylosing spondylitis, bursitis, burns, injuries.


25 Exemplifying the invention are the following compounds:



- 8 -



10 Illustrative of the pharmaceutically acceptable salts is the formula



wherein Ar is defined above and Z^+ is a pharmaceutically acceptable monovalent counterion. As is well appreciated by those of skill in the art, the pharmaceutically acceptable counterions include, aluminum, calcium, lithium, magnesium, potassium, sodium, barium, zinc, ammonium, or an amino acid such as glycine, alanine, valine, leucine, isoleucine, serine, threonine, aspartic acid, asparagine, glutamic acid, lysine, hydroxylysine, histidine, arginine, phenylalanine, tyrosine, tryptophan, thyroxine, cystine, cysteine, methionine, proline, hydroxyproline, ornithine, β -alanine, α -amino butyric acid, sarcosine, betaine, homoserine, and citrulline, or mono, di, or triC₁₋₆alkylamino.

Compounds of Formula I are useful for the relief of pain, fever and inflammation of a variety of conditions including rheumatic fever, symptoms associated with influenza or other viral infections, common cold, low back and neck pain, dysmenorrhea, headache, toothache, sprains and strains, myositis, neuralgia, synovitis, arthritis, including rheumatoid arthritis degenerative joint diseases (osteoarthritis), gout and ankylosing spondylitis, bursitis, burns,

- 9 -

injuries, following surgical and dental procedures. In addition, such compounds may inhibit cellular neoplastic transformations and metastatic tumor growth and hence can be used in the treatment of cancer.

Compounds of Formula I will also inhibit prostanoid-induced smooth muscle contraction by preventing the synthesis of contractile prostanoids and hence may be of use in the treatment of dysmenorrhea, premature labor and asthma. Compounds of formula I are also useful in the treatment of dementia, in particular, Alzheimer's disease.

By virtue of their high cyclooxygenase-2 (COX-2) activity and/or their specificity for cyclooxygenase-2 over cyclooxygenase-1 (COX-1), compounds of Formula I will prove useful as alternatives to conventional non-steroidal anti-inflammatory drugs (NSAID'S) particularly where such non-steroidal anti-inflammatory drugs may be contra-indicated such as in patients with peptic ulcers, gastritis, regional enteritis, ulcerative colitis, diverticulitis or with a recurrent history of gastrointestinal lesions; GI bleeding, coagulation disorders including anemia such as hypoprothrombinemia, haemophilia or other bleeding problems; kidney disease; those prior to surgery or taking anticoagulants.

Similarly, compounds of Formula I, will be useful as a partial or complete substitute for conventional NSAID'S in preparations wherein they are presently co-administered with other agents or ingredients. Thus in further aspects, the invention encompasses pharmaceutical compositions for treating cyclooxygenase-2 mediated diseases as defined above comprising a non-toxic therapeutically effective amount of compound of Formula I as defined above and one or more ingredients such as another pain reliever including acetaminophen or phenacetin; a potentiator including caffeine; an H2-antagonist, aluminum or magnesium hydroxide, simethicone, a decongestant including phenylephrine, phenylpropanolamine, pseudophedrine, oxymetazoline, ephinephrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine; an antitussive including codeine, hydrocodone, caramiphen, carbetapentane, or dextromethorphan; a diuretic; a sedating or non-

- 10 -

sedating antihistamine. In addition the invention encompasses a method of treating cyclooxygenase mediated diseases comprising:
administration to a patient in need of such treatment a non-toxic therapeutically effective amount of compound of Formula I, optionally
5 co-administered with one or more of such ingredients as listed immediately above.

Compounds of the present invention are inhibitors of cyclooxygenase-2 and are thereby useful in the treatment of cyclooxygenase-2 mediated diseases as enumerated above. This activity
10 is illustrated by their ability to selectively inhibit cyclooxygenase-2 over cyclooxygenase-1. Accordingly, in one assay, the ability of the compounds of this invention to treat cyclooxygenase mediated diseases can be demonstrated by measuring the amount of prostaglandin E₂ (PGE₂) synthesized in the presence of arachidonic acid,
15 cyclooxygenase-1 or cyclooxygenase-2 and a compound of formula I. The IC₅₀ values represent the concentration of inhibitor required to return PGE₂ synthesis to 50 % of that obtained as compared to the uninhibited control. Illustrating this aspect, we have found that the Compounds 1 through 26 are more than 100 times more effective in
20 inhibiting COX-2 than they are at inhibiting COX-1. In addition they have an IC₅₀ of 1 nM to 1 μ M. By way of comparison, Ibuprofen has an IC₅₀ for COX-2 of 1 μ M, and Indomethacin has an IC₅₀ for COX-2 of approximately 100nM.

For the treatment of any of these cyclooxygenase mediated
25 diseases compounds of Formula I may be administered orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal
30 injection or infusion techniques. In addition to the treatment of warm-blooded animals such as mice, rats, horses, cattle, sheep, dogs, cats, etc., the compounds of the invention are effective in the treatment of humans.

- 11 -

As indicated above, pharmaceutical compositions for treating cyclooxygenase-2 mediated diseases as defined may optionally include one or more ingredients as listed above.

5 The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical
10 compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically
15 acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or
20 acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also
25 be coated by the techniques described in the U.S. Patents 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release.

30 Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

- 12 -

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-
5 cellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic
10 alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan
15 monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the
20 active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a
25 palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending
30 agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

- 13 -

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of formula (I) may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

- 14 -

For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compounds of Formula (I) are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

5 Dosage levels of the order of from about 0.01 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions, or alternatively about 0.5 mg to about 7 g. per patient per day. For example, inflammation may be effectively treated by the administration of from about 0.01 to 50 mg of the
10 compound per kilogram of body weight per day, or alternatively about 0.5 mg to about 3.5 g per patient per day.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of
15 administration. For example, a formulation intended for the oral administration of humans may contain from 0.5 mg to 5 g of active agent compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Dosage unit forms will generally contain between
20 from about 1 mg to about 1000 mg of an active ingredient, typically 25 mg, 50 mg, 100 mg, 200 mg, 300 mg, 400 mg, 500 mg, 600 mg, 800 mg, or 1000 mg.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors
25 including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.


Compounds of the instant invention are conveniently
30 prepared using the procedures described in the methods below. Additional relevant chemistry is described in US 4,375,479, issued to Schroeder, et al., March 1, 1983 which is hereby incorporated by reference.

- 15 -

Method A

Indanol I is coupled with an appropriate alkylating agent and then nitrated to give the nitro ether II. Reduction, which can be performed by catalytic hydrogenation, followed by treatment with a sulfonylating agent such as an alkanesulfonyl chloride in the presence of a base like pyridine gives the sulfonamide III. Benzylic oxydation can be accomplished by a variety of reagents such as pyridinium chlorochromate in refluxing benzene to give the alkyl sulfonamido indanone IV.

Method A



- 16 -

Method B

Amino V is acetylated followed by bromination to give acetylamino bromo indane VI. Oxydation, which can be performed by many reagents such as chromium trioxide in acetic acid, followed by acid hydrolysis gives the amino indanone VII. Conversion of the amino group into a nitro group is done by treatment of the corresponding diazonium salt with sodium nitrite in the presence of copper. Protection of the carbonyl as a dioxolane provides the bromo nitro ketal VIII. Coupling with an appropriate nucleophile proceeds under basic conditions. Reduction of the nitro group with iron powder with concomitant hydrolysis of the ketal leads to the amino indanone IX. Monosulfonylation gives the alkanesulfonamido indanone X. In cases where monosulfonylation is impractical, the amino indanone IX is converted to the bissulfonamide followed by basic hydrolysis with aqueous hydroxide to give the alkanesulfonamido indanone X.


20

25

30

- 17 -


Method B



- 18 -

Method C

The bromo nitro indane VIII is converted to the ketone XI by cyanide displacement of the bromide followed by addition of an organometallic reagent such as a Grignard reagent onto the resulting nitrile and subsequent hydrolysis. Reduction and sulfonylation provides the sulfonamide XII which upon hydrolysis gives the alkanesulfonamido indanone XIII.



- 19 -


Method D

The bromonitroindane VIII is converted to the nitro thiophenol XIV either via disulfide formation and reductive cleavage or via thiolester displacement of the bromide followed by hydrolysis. This thiophenol is then alkylated and the nitro is reduced to give the amino thioether XV. Subsequent monosulfonylation gives the alkanesulfon-amido indanone XVI.

10

Method D


15



30

- 20 -

5



COMP.	X	R ¹	HETEROCYCLE	R ²	R ³
1	S	CH ₃	2-Thienyl	--	--
2	S	CH ₃	2-(1,3-)Diazinyl	4-CH ₃	--
3	S	CH ₃	2-Thiazolyl	--	--
4	S	CH ₃	2-(1,3,4-)Thiadiazolyl	--	--
5	O	CH ₃	3-Thienyl	--	--
6	S	CF ₃	2-thienyl	--	--
7	S	CH ₃	2-Thiazolyl	4-CH ₃	5-CH ₃
8	S	CH ₃	2-Imidazolyl	N-CH ₃	--
9	CH ₂	CF ₃	2-Thiazolyl	4-Cl	--
10	S	CH ₂ CH ₃	3-Furyl	2-CH ₃	--
11	S	CH(CH ₃) ₂	2-(1,3,4-)Thiadiazolyl	5-SCH ₃	--
12	S	CH ₃	2-(1,2,4-)Triazolyl	1-CH ₃	5-CH ₃
13	S	CH ₃	4-(1,2,3-)Triazolyl	1-CH ₃	--
14	O	CH ₃	4-Isotiazolyl	--	--
15	O	CF ₃	2-Thiazolyl	--	--
16	CH ₂	CH ₃	2-Oxazolyl	--	--
17	O	CH ₃	5-Thiazolyl	--	--
18	S	CH ₃	5-Isotiazolyl	3-CH ₃	--
19	O	CH ₃	3-Isotiazolyl	--	--
20	S	CH ₃	4-Isotiazolyl	3-CH ₃	--
21	O	CH ₃	4-(1,2,3-)Thiadiazolyl	--	--
22	S	CH ₃	4-(1,2,5-)Thiadiazolyl	--	--

- 21 -

COMP.	X	R ¹	HETEROCYCLE	R ²	R ³
23	S	CF ₃	3-Isioxazolyl	--	--
24	S	CH ₂ CH ₃	5-(1,2,3-)Thiadiazolyl	--	--
25	S	CH ₃	2-(1,3,4-)Oxadiazolyl	--	--
26	S	CF ₃	3-(1,2,4-)Thiadiazolyl	5-CH ₃	--
27	S	CH ₃	2-Thiazolyl	4-CH ₃	--
28	S	CH ₃	2-Thiazolyl	4-Et	--
29	S	CH ₃	2-Thiazolyl	4-i-Pr	--
30	S	CH ₃	2-Thiazolyl	4-t-Bu	--
31	S	CH ₃	2-Thiazolyl	4-C-Pr	--
32	S	CH ₃	2-Thiazolyl	4-vinyl	--

Using the methods A-D above, and the methodology described in Example 3, Compounds 1,2 and 4 were prepared.

The 4-substituted-2-mercaptothiazoles used in the synthesis Examples 5 and 6 may be prepared according to the method of Buchman (E.R. Buchman, A.O. Reims and H. Sargent, J. Org. Chem., 6, 764 (1941)).

EXAMPLE 1 (Compound 1)

5-Methanesulfonamido-6-(2-thiazolylthio)-1-indanone

Cal'd for Analysis
C, 49.54; H, 3.86; N, 4.13
Found: C, 49.37; H, 3.79; N, 3.97

EXAMPLE 2 (Compound 2)

5-Methanesulfonamido-6-(2-(4-methyl-1,3-diazinylthio))-1-indanone

High resolution mass spectrum (FAB):
MH⁺ found at 350.06330

- 22 -

calculated 350.06330

EXAMPLE 3 (Compound 3)5 5-Methanesulfonamido-6-(2-thiazolylthio)-1-indanone, sodium saltStep 1: 5-Acetylaminoindane

To a solution of 5-aminoindane (10.0 g, 7.5 meol) in CH₂Cl₂ (100 mL) was added dropwise acetic anhydride (9.2 g, 9.0 mmol) over a period of 15 min. After further stirring for 30 min, the mixture was quenched with 1 M aqueous NaOH (100 mL). The CH₂Cl₂ layer was separated, washed successively with 1 M aqueous HCl, brine, and was then dried over anhydrous MgSO₄ and concentrated *in vacuo*. Chromatography over silica gel, eluting with ethyl acetate:hexanes (1:1) afforded 12.2 g (85%) of the title compound as a light brown powder.

¹H NMR (CDCl₃): δ 7.44 (1H, s), 7.12 (3H, three overlapping s), 2.88 (4H, m), 2.15 (3H, s), 2.06 (2H, m).

20 Step 2: 5-Acetylamino-6-bromoindane

To a solution of 5-acetylaminoindane (53.0 g, 0.30 mmol) in glacial acetic acid (1 L) at 10°C was added dropwise over a period of 1 h a solution of bromine (19.0 mL, 0.37 meol). The mixture was further stirred at 10°C for 15 min, and was then diluted with water until no more precipitate formed. The precipitate was collected, washed with water and dried under vacuum to give 61 g (80%) of the title compound.

¹H NMR (CDCl₃): δ 8.14 (1H, s), 7.50 (1H, s), 7.38 (1H, s), 2.88 (4H, m), 2.20 (3H, s), 2.08 (2H, m).

30 Step 3: 5-Acetylamino-6-bromo-1-indanone

To a solution of 5-acetylamino-6-bromoindane (43.0 g, 0.17 mmol) in glacial acetic acid (400 mL) at 50-55°C was added dropwise a solution of chromium trioxide (70.0 g, 0.7 mmol) in 50%

- 23 -

aqueous acetic acid (400 mL) over a period of 30 min. After further stirring for 15 min, the mixture was cooled to 0°C and quenched with 2-propanol (100 mL). Solvent was removed *in vacuo*. The residue was diluted with water (1 L) and extracted with ethyl acetate (2 x 500 mL). The combined ethyl acetate layer was washed with 0.5 M aqueous NaOH (1 L), brine, dried over anhydrous MgSO₄ and concentrated to give 36 g (80%) of the title compound as a light brown solid which was contaminated with about 10% of 5-bromo-6-acetylamino-1-indanone. ¹H NMR (CDCl₃): δ 8.60 (1H, s), 7.98 (1H, s), 7.90 (1H, s), 3.10 (2H, t), 2.70 (2H, t), 2.30 (3H, s).

Step 4: 5-Amino-6-bromo-1-indanone

A mixture of 5-acetylamino-6-bromo-1-indanone (36.0 g, 0.13 mmol) and 6 M aqueous hydrochloric acid (800 mL) was refluxed for 1 h. The homogenous solution was then cooled to 0°C and adjusted to pH 8 with 10 M aqueous NaOH (~480 mL). The precipitate formed was collected, washed with water and dried under vacuum to afford 30.0 g (quantitative) of the title compound as a light brown powder. ¹H NMR (acetone-d₆): δ 7.65 (1H, s), 6.90 (1H, s), 5.80 (2H, br s), 2.95 (2H, t), 2.50 (2H, t).

Step 5: 5-Nitro-6-bromo-1-indanone

To a suspension of 5-amino-6-bromo-1-indanone (30.0 g, 0.13 mmol) in 20% aqueous fluoroboric acid (120 mL) at 0°C was added dropwise 4 M aqueous NaNO₂ (50 mL, 0.20 mmol) over a period of 30 min. The mixture was stirred for 30 min after completion of addition. The resulting foamy suspension was added portionwise to a vigorously stirred mixture of copper powder (40 g, 0.62 mmol) and sodium nitrite (120 g, 1.74 mmol) in water (240 mL) at room temperature over a period of 15 min. During the addition, excessive foaming was broken up by the addition of small amounts of diethyl ether. After further stirring for 30 min, the mixture was filtered through celite, washed with ethyl acetate (5 x 300 mL). The ethyl acetate layer was separated, washed with brine, dried over anhydrous

- 24 -

MgSO₄ and concentrated *in vacuo*. Chromatography over silica gel, eluting with CH₂Cl₂, yielded 17.5 g (51%) of the title compound as a pale yellow solid.

5 ¹H NMR (CDCl₃): δ 8.10 (1H, s), 7.85 (1H, s), 3.20 (2H, t), 2.85 (2H, t); mass spectrum (DCI, CH₄) m/e 256 (M⁺+H).

Step 6: 5-Nitro-6-bromo-1-indanone ethylene acetal

10 To a suspension of 5-nitro-6-bromo-1-indanone (11.0 g, 43 mmol) and bis(trimethylsilyloxy)ethane (22.0 mL, 90 mmol) in CH₂Cl₂ (90 mL) at room temperature was added trimethylsilyltrifluoromethanesulfonate (100 μL). The mixture was stirred for 2 h and the homogeneous solution was quenched with saturated aqueous NaHCO₃ (100 mL). The CH₂Cl₂ layer was separated, washed with brine, dried
15 over anhydrous MgSO₄ and concentrated *in vacuo*. Chromatography over silica gel, eluting with ethyl acetate:hexanes (2:5), furnished 10.2 g (79%) of the title compound as a pale yellow solid.

20 ¹H NMR (CDCl₃): δ 7.70 (1H, s), 7.68 (1H, s), 4.15 (4H, m), 2.98 (2H, t), 2.38 (2H, t)

Step 7: 5-Nitro-6-(2-thiazolylthio)-1-indanone ethylene acetal

25 The bromide from Step 6 (410 mg, 1.37 mmol) and 2-mercaptothiazole (200 mg, 1.7 mmol) were mixed together in pyridine (3 mL) at room temperature. Aqueous KOH (8 M, 213 μL, 1.7 mmol) was added and the resulting mixture was heated with an oil bath at 80°C for 3 h, then at 100°C for 1.5 h and finally 75°C for 16 h. After
30 cooling to room temperature the mixture was diluted and EtOAc and washed with water. The organic layer was dried over MgSO₄ and concentrated *in vacuo*. Purification by flash chromatography on silica gel (from 5% acetone in 2:1 CH₂Cl₂-hexanes to 10% acetone in CH₂Cl₂) yielded 280 mg of the title compound.

- 25 -

^1H NMR (CDCl_3): δ 8.50 (1H, s), 7.99 (1H, d), 7.58 (1H, d), 7.11 (1H, d), 4.05-3.90 (4H, m), 2.97 (2H, t), 2.32 (2H, t).

Step 8: 5-Amino-6-(2-thiazolylthio)-1-indanone

5 Iron powder (265 mg, 4.7 mmol), ammonium chloride (35 mg, 0.65 mmol) and the nitroindane from Step 7 (275 mg, 0.82 mmol) were mixed together in ethanol (5 mL) and water (2.5 mL). The mixture was refluxed for 1 h and then filtered through celite, washing the cake with EtOAc. The volatiles were removed *in vacuo* and the
10 residue was dissolved in EtOAc: CH_2Cl_2 (3:1), washed with brine and dried over MgSO_4 . Evaporation of the solvent left a residue that was purified by flash chromatography on silica gel (EtOAc:toluene: CH_2Cl_2 from 30:50:20 to 1:1:1) to give 150 mg of the title compound.

15 ^1H NMR (CDCl_3): δ 8.03 (1H, s), 7.67 (1H, d), 7.20 (1H, d), 6.78 (1H, d), 5.05 (2H, br s), 3.05 (2H, m), 2.67 (2H, m).

Step 9: 5-Bis(methanesulfonyl)amino-6-(2-thiazolylthio)-1-indanone

20 Methanesulfonyl chloride (710 μL , 9.15 mmol) and pyridine (740 μL , 9.15 mmol) were added to a solution of the aminoindanone from Step 8 in CH_2Cl_2 (5 mL). After stirring overnight at room temperature, Et_3N (400 μL , 2.8 mmol) and methanesulfonyl chloride (220 μL , 2.8 mmol) were added. The
25 mixture was stirred 4 h at room temperature. Water and CH_2Cl_2 were added, the aqueous phase was extracted with CH_2Cl_2 and the combined organic phase was washed with 0.1 M aqueous citric acid, dried over MgSO_4 and concentrated *in vacuo*. The crude residue was purified by flash chromatography on silica gel (20% EtOAc in CHCl_3) to give 175
30 mg of the title compound.

^1H NMR (CDCl_3): δ 7.96 (1H, s), 7.79 (1H, d), 7.58 (1H, s), 7.39 (1H, d), 3.55 (6H, s), 3.21 (2H, m), 2.77 (2H, m).

- 26 -

Step 10: 5-Methanesulfonamido-6-(2-thiazolylthio)-1-indanone)

The bismethanesulfonamide from Step 9 (175 mg, 0.42 mmol) was dissolved in THF (4 mL) and methanol (500 μ L) at room temperature. Aqueous NaOH (2.0 M, 630 μ L, 1.25 mmol) was added and the mixture stirred for 30 min. Saturated aqueous ammonium chloride was added and the mixture was extracted twice with EtOAc. The organic layer was dried over MgSO₄ and concentrated *in vacuo*. The crude was purified by flash chromatography (EtOAc:CH₂Cl₂:2-propanol, from 15:83:2 to 25:70:5) to give 96 mg of the title compound.

¹H NMR (CDCl₃): δ 8.91 (1H, br s), 8.11 (1H, s), 7.88 (1H, s), 7.71 (1H, d), 7.32 (1H, d), 3.18 (2H, m), 3.07 (3H, s), 2.72 (2H, m).

Step 11: 5-Methanesulfonamido-6-(2-thiazolylthio)-1-indanone, sodium salt

The methanesulfonamide of Step 10 (96 mg, 0.28 mmol) was dissolved in ethanol (1 mL). Aqueous NaOH (1.0 M, 280 μ L, 0.28 mmol) was added followed by water (5 mL). The mixture was concentrated *in vacuo* and the aqueous residue was freeze-dried overnight to give 100 mg of the title compound.

High Resolution Mass (FAB)

MH⁺ found at 362.99078


MH⁺ calculated at 362.9907775

EXAMPLE 4 (Compound 4)5-Methanesulfonamido-6-(2-(5-methyl-1,3,4-thiadiazoylthio))-1-indanone


[M+Na]⁺=378

EXAMPLE 5 (Compound 27)

- 27 -

5-Methanesulfonamido-6-(4-methyl-2-thiazolylthio)-1-indanone

Step 1: 5-Nitro-6-(4-methyl-2-thiazolylthio)-1-indanone ethylene ketal




The compound was prepared in the same manner as Step 7 Example 3, but using 4-methyl-2-mercaptothiazole (Buchman, et al., J. Org. Chem., 1941, 6, 764).

¹H NMR (CDCl₃): δ 8.02 (1H, s), 7.25 (1H, s), 7.10 (1H, s), 4.10-3.90 (4H, m), 3.00 (2H, t), 2.55 (3H, s), 2.35 (2H, t).

Step 2: 5-Amino-6-(4-methyl-2-thiazolylthio)-1-indanone

- 28 -



5


The compound was prepared in the same manner as Step 8
Example 3.

10

¹H NMR (CDCl₃): δ 8.00 (1H, s), 6.75 (1H, s), 6.72 (1H, s), 4.30 (2H, brs), 3.05 (2H, t), 2.65 (2H, t), 2.40 (3H, s).

15

Step 3: 5-Bis(methanesulfonyl)amino-6-(4-methyl-2-thiazolylthio)-1-indanone



20

25


The compound was prepared in the same manner as Step 9
Example 3.

30

¹H NMR (CDCl₃): δ 7.94 (1H, s), 7.56 (1H, s), 6.95 (1H, s), 3.55 (6H, s), 3.20 (2H, t), 2.75 (2H, t), 2.45 (3H, s).

Step 4: 5-Methanesulfonamido-6-(4-methyl-2-thiazolylthio)-1-indanone

- 29 -



5


The compound was prepared in the same manner as Step 10
Example 3.

10

¹H NMR (CDCl₃): δ 9.12 (1H, brs), 8.05 (1H, s), 7.88 (1H, s), 6.85 (1H, s), 3.20 (2H, t), 3.10 (3H, s), 2.70 (2H, t), 2.42 (3H, s).

EXAMPLE 6 (Compound 28)

15




20

5-Methanesulfonamido-6-(4-ethyl-2-thiazolythio)-1-indanone

25

Step 1: 5-Nitro-6-(4-ethyl-2-thiazolythio)-1-indanone ethyl ketal




30

- 30 -

The compound was prepared in the same manner as Step 7 Example 3, but using 4-ethyl-2-mercaptothiazole (Buchman, et al., J. Org. Chem., 1941, 6, 764).

¹H NMR (CDCl₃): δ 8.03 (1H, s), 7.16 (1H, s), 7.10 (1H, s), 3.99 (2H, m) 3.91 (2H, m), 2.96 (2H, t), 2.85 (2H, q), 2.30 (2H, t), 1.32 (3H, t).


Step 2: 5-Amino-6-(4-ethyl-2-thiazolythio)-1-indanone



The compound was prepared in the same manner as Step 8 Example 3.

¹H NMR (CDCl₃): δ 8.00 (1H, s), 6.75 (1H, s), 6.71 (1H, s), 5.05 (2H, brs), 3.04 (2H, m) 2.73 (2H, q), 2.65 (2H, m), 1.25 (3H, t).

Step 3: 5-Bis(methanesulfonyl)amino-6-(4-ethyl-2-thiazolythio)-1-indanone



- 31 -


The compound was prepared in the same manner as Step 9
Example 3.

5 ^1H NMR (CDCl_3): δ 7.93 (1H, s), 7.52 (1H, s), 6.91 (1H, s), 3.53 (6H, s), 3.18 (2H, m), 2.75 (4H, m), 1.25 (3H, t).

Step 4: 5-Methanesulfonamido-6-(4-ethyl-2-thiazolythio)-1-
indanone

10

15



The compound was prepared in the same manner as Step 10
Example 3.

20

^1H NMR (CDCl_3): δ 9.42 (1H, brs), 8.08 (1H, s), 7.85 (1H, s), 6.83 (1H, s), 3.15 (2H, m), 3.05 (3H, s), 2.75 (4H, m), 1.25 (3H, t).


25

30

- 32 -

WHAT IS CLAIMED IS:

1. A compound of formula I



I

or a pharmaceutically acceptable salt thereof wherein:

R¹ is selected from the group consisting of

- 15
- (a) C₁-7alkyl, and
 - (b) mono-, di-, tri-, tetra- and per-substituted C₁-7alkyl, wherein the substituent is fluoro;

X is O, S, or -CH₂-;

Ar is a mono or disubstituted aromatic ring, said ring selected from

- 20
- (a) a ring of 5 atoms containing one O, S or N atom and optionally, 1, 2 or 3 additional N atoms; and
 - (b) a ring of 6 atoms containing 2, 3 or 4 nitrogen atoms;

wherein the substituents are independently selected from

- 25
- (a) hydrogen,
 - (b) C₁-7alkyl,
 - (c) halogen, including F, Cl, Br, and I,
 - (d) OCH₃,
 - (e) SCH₃,
 - 30 (f) CF₃,
 - (g) COCH₃ and
 - (h) S(O)₂CH₃.

- 33 -

2. A compound according to Claim 1 wherein:

Ar is a mono or di-substituted aromatic group wherein the aromatic group is selected from the group consisting of

- (1) furanyl,
- 5 (2) diazinyl, triazinyl and tetrazinyl,
- (3) imidazolyl,
- (4) isooxazolyl,
- (5) isothiazolyl,
- (6) oxadiazolyl,
- 10 (7) oxazolyl,
- (8) pyrazolyl,
- (9) pyrrolyl,
- (10) thiadiazolyl,
- (11) thiazolyl,
- 15 (12) thienyl,
- (13) triazolyl, and
- (14) tetrazolyl,

wherein the substituents are independently selected from

- (a) hydrogen,
- 20 (b) C₁₋₇alkyl,
- (c) halogen,
- (d) OCH₃,
- (e) SCH₃,
- (f) CF₃,
- 25 (g) COCH₃, and
- (h) S(O)₂CH₃.

3. A compound according to Claim 2 wherein

R¹ is selected from the group consisting of

- 30 (a) methyl, ethyl, propyl and butyl,
- (b) mono, di or tri substituted methyl, ethyl, propyl or butyl, wherein the substituent is fluoro;

- 34 -

X is O,

Ar is a mono or di substituted aromatic group wherein the aromatic group is selected from the group consisting of

- (1) diazinyl,
- (2) isooxazolyl,
- (3) isothiazolyl,
- (4) oxadiazolyl,
- (5) oxazolyl,
- (6) thiadiazolyl, and
- (7) thiazolyl,

wherein the substituents are independently selected from

- (a) hydrogen,
- (b) C₁-3alkyl,
- (c) F or Cl,
- (d) OCH₃,
- (e) SCH₃, and
- (f) CF₃.

4. A compound according to Claim 3 wherein
- R¹ is selected from the group consisting of
- (a) methyl, ethyl or propyl, and
 - (b) mono, di or tri substituted methyl, ethyl or propyl, wherein the substituent is fluoro;

X is O;

Ar is a mono or di substituted aromatic group wherein the aromatic group is selected from the group consisting of

- (1) 3-isooxazolyl,
- (2) 4-isooxazolyl,
- (3) 5-isooxazolyl,
- (4) 3-isothiazolyl,
- (5) 4-isothiazolyl,
- (6) 5-isothiazolyl,
- (7) 2-oxazolyl,
- (8) 4-oxazolyl,

- 35 -

- 5 (9) 5-oxazolyl,
(10) 2-thiazolyl,
(11) 4-thiazolyl,
(12) 5-thiazolyl,
(13) 1,2,3-thiadiazol-4-yl,
(14) 1,2,3-thiadiazol-5-yl,
(15) 1,2,4-thiadiazol-3-yl,
(16) 1,2,4-thiadiazol-5-yl,
(17) 1,3,4-thiadiazol-2-yl,
10 (18) 1,2,5-thiadiazol-3-yl,
(19) 1,2,3-oxadiazol-4-yl,
(20) 1,2,3-oxadiazol-5-yl,
(21) 1,2,4-oxadiazol-3-yl,
(22) 1,2,4-oxadiazol-5-yl,
15 (23) 1,3,4-oxadiazol-2-yl,
(24) 1,2,5-oxadiazol-3-yl,
(25) 1,2-diazinyl,
(26) 1,3-diazinyl, and
(27) 1,4-diazinyl,
20 wherein the substituents are independently selected from
(a) hydrogen,
(b) methyl, ethyl or propyl,
(c) F or Cl, and
(d) OCH₃.

25

5. A compound according to Claim 3 wherein

R¹ is methyl

X is O;

Ar is a mono or di substituted aromatic group wherein the aromatic
30 group is selected from the group consisting of

- (1) 3-isothiazolyl,
(2) 4-isothiazolyl,
(3) 5-isothiazolyl,
(4) 2-oxazolyl,

- 36 -

- 5
- (5) 4-oxazolyl,
 - (6) 5-oxazolyl,
 - (7) 2-thiazolyl,
 - (8) 4-thiazolyl,
 - (9) 5-thiazolyl,
 - (10) 1,2-diazinyl,
 - (11) 1,3-diazinyl, and
 - (12) 1,4-diazinyl,

wherein the substituents are independently selected from

- 10
- (a) hydrogen,
 - (b) methyl, ethyl or propyl,
 - (c) F or Cl, and
 - (d) OCH₃.

- 15
6. A compound according to Claim 2 wherein R¹ is selected from the group consisting of
- (a) methyl, ethyl, propyl and butyl,
 - (b) mono, di or tri substituted methyl, ethyl, propyl or butyl, wherein the substituent is Fluoro;

20 X is CH₂,

Ar is a mono or di substituted aromatic group wherein the aromatic group is selected from the group consisting of

- 25
- (1) diazinyl,
 - (2) isooxazolyl,
 - (3) isothiazolyl,
 - (4) oxadiazolyl,
 - (5) oxazolyl,
 - (6) thiadiazolyl, and
 - (7) thiazolyl,

30 wherein the substituents are independently selected from

- (a) hydrogen,
- (b) C₁-3alkyl,
- (c) F or Cl,
- (d) OCH₃,

- 37 -

- (e) SCH₃, and
- (f) CF₃.

7. A compound according to Claim 6 wherein
- 5 R¹ is selected from the group consisting of
- (a) methyl, ethyl or propyl, and
 - (b) mono, di or tri substituted methyl, ethyl or propyl, wherein the substituent is fluoro;

X is CH₂;

- 10 Ar is a mono or di substituted aromatic group wherein the aromatic group is selected from the group consisting of

- (1) 3-isooxazolyl,
- (2) 4-isooxazolyl,
- (3) 5-isooxazolyl,
- 15 (4) 3-isothiazolyl,
- (5) 4-isothiazolyl,
- (6) 5-isothiazolyl,
- (7) 2-oxazolyl,
- (8) 4-oxazolyl,
- 20 (9) 5-oxazolyl,
- (10) 2-thiazolyl,
- (11) 4-thiazolyl,
- (12) 5-thiazolyl,
- (13) 1,2,3-thiadiazol-4-yl,
- 25 (14) 1,2,3-thiadiazol-5-yl,
- (15) 1,2,4-thiadiazol-3-yl,
- (16) 1,2,4-thiadiazol-5-yl,
- (17) 1,3,4-thiadiazol-2-yl,
- (18) 1,2,5-thiadiazol-3-yl,
- 30 (19) 1,2,3-oxadiazol-4-yl,
- (20) 1,2,3-oxadiazol-5-yl,
- (21) 1,2,4-oxadiazol-3-yl,
- (22) 1,2,4-oxadiazol-5-yl,
- (23) 1,3,4-oxadiazol-2-yl,

- 38 -

- (24) 1,2,5-oxadiazol-3-yl,
- (25) 1,2-diazinyl,
- (26) 1,3-diazinyl, and
- (27) 1,4-diazinyl,

5 wherein the substituents are independently selected from

- (a) hydrogen,
- (b) methyl, ethyl or propyl,
- (c) F or Cl, and
- (d) OCH₃.

10

8. A compound according to Claim 7 wherein

R¹ is methyl

X is CH₂ ;

Ar is a mono or di substituted aromatic group wherein the aromatic
15 group is selected from the group consisting of

15

- (1) 3-isothiazolyl,
- (2) 4-isothiazolyl,
- (3) 5-isothiazolyl,
- (4) 2-oxazolyl,
- (5) 4-oxazolyl,
- (6) 5-oxazolyl,
- (7) 2-thiazolyl,
- (8) 4-thiazolyl,
- (9) 5-thiazolyl,
- (10) 1,2-diazinyl,
- (11) 1,3-diazinyl, and
- (12) 1,4-diazinyl,

20

25

wherein the substituents are independently selected from

- (a) hydrogen,
- (b) methyl or ethyl,
- (c) F or Cl, and
- (d) OCH₃.

30

9. A compound according to Claim 2 wherein

- 39 -

R¹ is selected from the group consisting of

- (a) methyl, ethyl or propyl, and
- (b) mono, di or tri substituted methyl, ethyl, propyl or butyl, wherein the substituent is Fluoro;

5 X is S;

Ar is a mono or di substituted aromatic group wherein the aromatic group is selected from the group consisting of

- (1) diazinyl,
- (2) isooxazolyl,
- 10 (3) isothiazolyl,
- (4) oxadiazolyl,
- (5) oxazolyl,
- (6) thiadiazolyl, and
- (7) thiazolyl,

15 wherein the substituents are independently selected from

- (a) hydrogen,
- (b) C₁₋₃alkyl,
- (c) F or Cl,
- (d) OCH₃,
- 20 (e) SCH₃, and
- (f) CF₃.

10. A compound according to Claim 9 wherein

R¹ is selected from the group consisting of

- 25 (a) methyl, ethyl or propyl, and
- (b) mono, di or tri substituted methyl, ethyl or propyl, wherein the substituent is fluoro;

X is S;

Ar is a mono or di substituted aromatic group wherein the aromatic group is selected from the group consisting of

- 30 (1) 3-isooxazolyl,
- (2) 4-isooxazolyl,
- (3) 5-isooxazolyl,
- (4) 3-isothiazolyl,

- 40 -

- 5
10
15
20
- (5) 4-isothiazolyl,
 - (6) 5-isothiazolyl,
 - (7) 2-oxazolyl,
 - (8) 4-oxazolyl,
 - (9) 5-oxazolyl,
 - (10) 2-thiazolyl,
 - (11) 4-thiazolyl,
 - (12) 5-thiazolyl,
 - (13) 1,2,3-thiadiazol-4-yl,
 - (14) 1,2,3-thiadiazol-5-yl,
 - (15) 1,2,4-thiadiazol-3-yl,
 - (16) 1,2,4-thiadiazol-5-yl,
 - (17) 1,3,4-thiadiazol-2-yl,
 - (18) 1,2,5-thiadiazol-3-yl,
 - (19) 1,2,3-oxadiazol-4-yl,
 - (20) 1,2,3-oxadiazol-5-yl,
 - (21) 1,2,4-oxadiazol-3-yl,
 - (22) 1,2,4-oxadiazol-5-yl,
 - (23) 1,3,4-oxadiazol-2-yl,
 - (24) 1,2,5-oxadiazol-3-yl,
 - (25) 1,2-diazinyl,
 - (26) 1,3-diazinyl, and
 - (27) 1,4-diazinyl,

wherein the substituents are independently selected from

- 25
- (a) hydrogen,
 - (b) methyl, ethyl or propyl,
 - (c) F or Cl, and
 - (d) OCH₃.

- 30
11. A compound according to Claim 10 wherein
R¹ is methyl,
X is S;
Ar is a mono or di substituted aromatic group wherein the aromatic
group is selected from the group consisting of


- 41 -

- 5
10
- (1) 3-isothiazolyl,
 - (2) 4-isothiazolyl,
 - (3) 5-isothiazolyl,
 - (4) 2-oxazolyl,
 - (5) 4-oxazolyl,
 - (6) 5-oxazolyl,
 - (7) 2-thiazolyl,
 - (8) 4-thiazolyl,
 - (9) 5-thiazolyl,
 - (10) 1,2-diazinyl,
 - (11) 1,3-diazinyl, and
 - (12) 1,4-diazinyl,

wherein the substituents are independently selected from

- 15
- (a) hydrogen,
 - (b) methyl or ethyl,
 - (c) F or Cl, and
 - (d) OCH₃.

20 12. A compound of formula



Z is pharmaceutically acceptable monovalent counterion,

Ar is a mono or di substituted aromatic group wherein the aromatic
30 group is selected from the group consisting of

- (1) 3-isothiazolyl,
- (2) 4-isothiazolyl,
- (3) 5-isothiazolyl,
- (4) 2-oxazolyl,



- 42 -

- (5) 4-oxazolyl,
(6) 5-oxazolyl,
(7) 2-thiazolyl,
(8) 4-thiazolyl,
(9) 5-thiazolyl,
(10) 1,2-diazinyl,
(11) 1,3-diazinyl, and
(12) 1,4-diazinyl,



wherein the substituents are independently selected from

- (a) hydrogen,
(b) methyl or ethyl,
(c) F or Cl, and
(d) OCH₃.

13. A compound according to Claim 12 selected from the group consisting of



and



and

and

- 43 -

or a pharmaceutically acceptable salt thereof.

5 14. A pharmaceutical compositions of inhibiting cyclooxygenase comprising a pharmaceutically acceptable carrier and a non-toxic therapeutically effective amount of compound according to Claim 1.

10 15. A pharmaceutical compositions of inhibiting cyclooxygenase-2 comprising a pharmaceutically acceptable carrier and a non-toxic therapeutically effective amount of compound according to Claim 1.

15 16. A method of inhibiting cyclooxygenase comprising: administration to a patient in need of such inhibition of a non-toxic therapeutically effective amount of a compound according to Claim 1.

20 17. A method of inhibiting cyclooxygenase-2 comprising: administration to a patient in need of such treatment of a non-toxic therapeutically effective amount of a compound according to Claim 1.

25 18. A pharmaceutical composition for the treatment of cyclooxygenase-2 mediated disease comprising a non-toxic therapeutically effective amount of compound according to Claim 1 and at least one or more ingredients selected from a pain reliever including acetaminophen or phenacetin; a potentiator including caffeine; an H₂-antagonist, aluminum or magnesium hydroxide, simethicone, a decongestant including phenylephrine, phenylpropanolamine, pseudopheorine, oxymetazoline, ephinephrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine; an
30 antitussive including codeine, hydrocodone, caramiphen, carbetapentane, or dexamethorphan; a diuretic; and a sedating or non-sedating antihistamine.

- 44 -

19. A pharmaceutically acceptable salt of a compound of formula (I) as defined in claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11.

20. A cyclooxygenase-2 inhibiting pharmaceutical composition comprising a non-toxic cyclooxygenase-2 inhibiting amount of a
5 compound of formula (I), or a pharmaceutically acceptable salt thereof, as defined in claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11, in association with a pharmaceutically acceptable carrier.

21. A cyclooxygenase-2 inhibiting pharmaceutical composition comprising a non-toxic cyclooxygenase-2 inhibiting amount of a
10 compound or salt of claim 12 or 13, in association with a pharmaceutically acceptable carrier.

22. A compound of formula (I), or a pharmaceutically acceptable salt thereof, as defined in claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11, for use in the treatment of cyclooxygenase-2 mediated disease.

15 23. Use of a compound or salt of claim 12 or 13 in the manufacture of a medicament for use in the treatment of cyclooxygenase-2 mediated disease.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA 94/00135

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C07D277/36 C07D239/38 C07D285/12 A61K31/425 A61K31/41
A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 009 554 (SCHERING AG) 16 April 1980 see page 4, line 1 - page 4, line 12; claims 1-75 ---	1-23
Y	EP,A,0 056 956 (SCHERING AG) 4 August 1982 see page 4, line 10 - page 4, line 13; claims 1-12 ---	1-23
Y	EP,A,0 140 684 (MERCK FROSST CANADA INC.) 8 May 1985 see page 4, line 26 - page 5, line 33; claims 1-30 --- -/--	1-23



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 June 1994

Date of mailing of the international search report

- 6. 07. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Herz, C

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, vol. 98, no. 9, 28 February 1983, Columbus, Ohio, US; abstract no. 65117m, C. RUFER ET AL. 'Nonsteroidal antiinflammatories. XII. Mode of action of antiinflammatory methane sulfonanilides' page 16 ; see abstract & Biochem. Pharmacol. 31 (22) 3591-6 (1982) ----	1-23
Y	CHEMICAL ABSTRACTS, vol. 107, no. 21, 23 November 1987, Columbus, Ohio, US; abstract no. 190553f, I. BOETTCHER ET AL. 'A sulfonamido-indanone derivative CGP 28237 (ZK 34228), a novel non-steroidal anti-inflammatory agent without gastro-intestinal ulcerogenicity in rats' page 37 ; see abstract & Drugs Exp. Clin. Res. 13 (5) 237-45 (1987) ----	1-23
Y	CHEMICAL ABSTRACTS, vol. 110, no. 15, 10 April 1989, Columbus, Ohio, US; abstract no. 128249v, I. WIESENBERG-BOETTCHER ET AL. 'The pharmacological profile of CGP 28238, a highly potent anti-inflammatory compound' page 39 ; see abstract & Agents Actions 26 (1-2) 240-2 (1989) -----	1-23

INTERNATIONAL SEARCH REPORT

information on patent family members

International application No.

PCT/CA 94/00135

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0009554	16-04-80	DE-A- 2833202	14-02-80
		DE-A- 2923937	08-01-81
		AT-T- 3148	15-05-83
		AU-B- 532405	29-09-83
		AU-A- 4932579	07-02-80
		CA-A- 1124724	01-06-82
		FR-A- 2433512	14-03-80
		GB-A- 2025973	30-01-80
		JP-C- 1463029	28-10-88
		JP-A- 55020777	14-02-80
		JP-B- 63010696	08-03-88
		SU-A- 965352	07-10-82
		US-A- 4244960	13-01-81
EP-A-0056956	04-08-82	DE-A- 3103372	02-09-82
		AU-B- 555252	18-09-86
		AU-A- 7988782	05-08-82
		CA-A- 1175443	02-10-84
		GB-A- 2092144	11-08-82
		JP-B- 1050698	31-10-89
		JP-C- 1562757	12-06-90
		JP-A- 57176948	30-10-82
		US-A- 4375479	01-03-83
EP-A-0140684	08-05-85	CA-A- 1254578	23-05-89
		JP-A- 60115544	22-06-85
		US-A- 4983628	08-01-91
		US-A- 4683325	28-07-87

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.